

# Deep Throat Oral Anesthetic Spray

## **Drug Facts**

<b>Active ingredient</b>	<b>Purpose</b>
Benzocaine 5 % .....	Oral Anesthetic
Glycerin 33 % .....	Oral Demulcent

### **Uses** For Temporary Relief of:

- occasional minor irritation, pain, sore mouth and sore throat
- pain due to canker sores, minor irritation or injury of the mouth and gums
- minor discomfort and protection of irritated areas in sore mouth and sore throat

### **Warnings**

- Allergy alert: Do not use this product if you have a history of allergy to local anesthetics such as procaine, butacaine, benzocaine or any other 'caine' anesthetics.
- Sore Throat Warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, swelling, nausea, or vomiting, consult a doctor promptly.
- Stop use and ask a doctor if sore mouth symptoms do not improve in 7 days, or if irritation, pain or redness persists or worsens.
- If pregnant or breast feeding, ask a health professional before use.
- **Keep this and all drugs out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.
- Do not exceed recommended dosage.
- Flammable. Do not use near fire, flame or heat.
- When using this product, do not get into eyes. If contact occurs, rinse eyes thoroughly with water. If irritation persists, consult a doctor.

### **Directions**

- Shake well before use.
- Adults: spray into throat or onto affected area with only ONE spray per use. Use up to 4 times daily or as directed by a doctor or dentist.

### **Other Information**

- Store at 20-25°C (68-77°F)
- Do not freeze.

### **Inactive Ingredients**

Alcohol Denat., Flavor (Aroma), PEG-40 Hydrogenated Castor Oil, PVP, Sucralose, Water (Aqua). May Contain: Caramel, Blue 1 (CI 42090), Red 40 (CI16035), Yellow 5 (CI 19140)